

Improving reusability of Atmospheric Model Data by adapting metadata standards and by providing specific data quality information

D. Neumann¹, **K. Peters**¹, A. Ganske², J. Mülmenstädt³, V. Voss⁴, D. Grawe⁴, H. Höck¹, S. Kindermann¹, A. Kraft², A. Lammert¹, B. Leitl⁴, J. Quaas³, H. Schlünzen⁴, H. Thiemann¹











GEFÖRDERT VOM



Motivation



- data publication numbers (incl. DataCite DOI's) are increasing, however...
- data reusability often difficult
 - wide variety of file formats
 - incomplete metadata
 - insufficient quality information

Motivation for AtMoDat: improve the reusability of atmospheric model data

- affects entire scientific communities
 - data users: reusable data difficult to identify
 - data producers and repositories: less reuse of data, reduced publicity
 - o funding institutions: project outcomes not reusable



Tasks in AtMoDat



support small subdisciplines in developing feasible standardization for their model output:



small MIPs



quality indicator for
DataCite DOI metadata
schema including
discipline-specific
extensions^[1,2]





Provide extended **DataCite DOI** with **quality indicator** to the atmospheric research community



Quality Indicator



- extension DataCite DOI metadata schema by "quality assessment" or "quality level" [1]
- quality criteria:
 - non-mandatory DataCite metadata filled
 - o discipline-specific metadata [2]
 - o open file format
 - quality control of metadata by data repository

Example for quality levels (atmospheric model data)

- 0. no quality control
- optional DataCite metadata provided
- 2. netCDF-CF (+ level 1)
- 3. discipline specific metadata and quality control (+ level 2)

The end of talk – the beginning of your involvement



Take part in our **survey** for urban atmospheric modelers:

Visit us at https://www.atmodat.de

Thank you for your attention!



uhh.de/orm-survey

GEFÖRDERT VOM









