

## Climate projections for West Africa: highly needed but uncertain

With climate change being one of the most severe challenges to rural Africa in the 21st century, West Africa is facing an urgent need to develop effective adaptation and mitigation measures to protect its constantly growing population. WASCAL (West African Science Service Center on Climate Change and Adapted Land Use, <http://www.wascal.org>) is a large-scale research-focused program designed to enhance the resilience of human and environmental systems to climate change and increased variability. An integral part of its climate services is the provisioning of a set of high resolution, ensemble-based regional climate projections for continental West Africa.

## II. Towards a regional climate modeling system for West Africa

An optimal model configuration of each RCM is required to reduce biases and reproduce the observed annual cycle of the WA monsoon precipitation. Klein et al. (2015) investigated the performance of 29 WRF model configurations with respect to monsoon dynamics, precipitation patterns, temperature and wind systems (Fig. 3). We evaluated their most promising candidate configurations using MPI-ESM MR data to derive a reference configuration (Table 1). For the Cosmo-CLM runs, we adopt the CORDEX WA setup identified by Panitz et al. (2012), while for the RegCM runs, the Sylla et al. (2010) setup is used.

Table 1. WRF setup

Microphysics	WSM5
Radiation SW/LW	RRTMG
Cumulus	GD
Boundary layer	ACM2
Surface layer	MM5
Land surface	Noah





Fig 3. (left) Box plots of precipitation bias wrt. ensemble mean. CU mostly affects single events, MP the amount of rainfall and PBL the position of the rain band. (top) Hovmöller diagrams of 1999 daily rain (Klein et al., 2015).

## I. The WASCAL regional climate projections: experiment design

To determine the uncertainty in the regional climate simulations stemming from the driving global circulation model and the limited area model itself, we adopt an ensemble approach (c.f., ENSEMBLES, CORDEX), where the multi-model mean and the spread of the ensemble describe the most likely projection and the uncertainty of it. We combine three global circulation models (GCM) and three regional climate models (RCM) for the future emission scenario RCP4.5. We also conduct historical runs and control runs using re-analysis data (Fig. 1).





Fig 1. WASCAL experiments: historical (1979-2005) and RCP4.5 (2006-2100) runs

## II. The WASCAL WRF simulations: model and domain setup

The governing meteorological feature over West Africa is the summer monsoon. This annual change in large-scale circulation and pressure systems plays a pivotal role in people's life as it provides the majority of the annual precipitation and the basis for agriculture in the region. As shown in a recent study by Brown and Sylla (2012), a large domain centered over West Africa is required to capture these processes and to generate the mesoscale convective cells. To provide added value over existing projections, we use a high spatial resolution of 12km in a 5:1 nesting setup. We use spectral nudging to keep the outer domain aligned with the driving model (Fig 2).




A large set of output variables on 24 pressure levels is provided every 3hr model time (see the WASCAL website for a complete list). Initial tests of two different methods to bias-correct the forcing GCM data led to mixed results (Heinzeller et al., 2014). Accordingly, no bias correction is applied to the forcing data in our simulations.

Fig. 2. Inner domain d02, 500x330x40 grid points, terrain height in m.


## IV. Model validation: historical and control runs

WRF runs at 12km resolution, driven by ERA-Interim re-analysis (R), MPI-ESM MR (E) and GFDL-ESM2M (G) for the period 1980-2010. Additionally, CCLM runs at 12km, WRF runs at 60km (outer domain), observational data from UDEL and two CORDEX runs using RCA4 are displayed (Colins et al., 2012). Observed mean surface temperatures are matched best by the 12km WRF/CCLM-R control runs and the 12km WRF-E run. GFDL-driven models show a distinct cold bias. Monthly rainfall over WA is closest to UDEL for the 12km control runs and the GFDL-driven historical runs. The WRF-R/E runs over-predict rainfall in Central Africa. Offshore, GFDL-driven models produce extreme precipitation with effects on Guinea.



## V. First projections: a warmer and wetter future?

First results for the 12km WRF-E runs show a warming signal of 2°C towards the end of the 21st century, in accordance with the CORDEX data. WRF-E indicates an increase in precip for WA, while CORDEX-G predicts a weaker trend and CORDEX-E a drier future for some areas. This illustrates the uncertainty in climate projections.



## References

- Brown, N. A. K., Sylla, M. B.: Regional Climate Model Sensitivity to Domain Size for the Simulation of the West African Summer Monsoon Rainfall. International Journal of Geophysics, Article ID 625831, 2012
- Nikulin, G.: Regional Climate Modeling Results and Ensemble using RCA4. RICCAR, The Fifth Expert Group Meeting, held December 11–12 2013 in Amman, Jordan
- Heinzeller, D., Klein, C., Kunstmann, H.: Towards a climate modeling system for West Africa - Sensitivity studies and input bias correction for WRF. 15th Annual WRF Users' Workshop, held June 24–26 2014 in Boulder, CO, USA
- Klein, C., Heinzeller, D., Bliefernicht, J., Kunstmann, H.: Variability of West African monsoon patterns generated by a WRF multi-physics ensemble. Climate Dynamics, DOI: 10.1007/s00382-015-2505-5, 2015
- Panitz, H.-J., Dosio, A., Büchner, M., Lüthi, D., Keuler, K.: COSMO-CLM (CCLM) climate simulations over CORDEX-Africa domain: analysis of the ERA-Interim driven simulations at 0.44° and 0.22° resolution. Climate Dynamics, 42, 3015–3038, 2014
- Sylla, M. B., Dell'Aquila, A., Ruti, P. M., Giorgi, F.: Simulation of the Intraseasonal and the Interannual Variability of Rainfall over West Africa with a Regional Climate Model (RegCM3) during the Monsoon Period. Int J of Climatology, 30, 1865–1883, 2010
- Jones, C., Nikulin, G., Kjellström, E., and 8 co-authors: Africa-CORDEX simulations, Canadian Meteorological and Oceanographic Society (CMOS) Congress 2012, held May 29–June 1 2012 in Montréal, Canada

## So, where is my added value?

Our historical runs indicate that the 12km models reproduce the annual cycle in temperature and rainfall better than existing CORDEX runs, and provide more accurate results for the sub-saharan regions. This comparison is hampered by the fact that up to now, no CORDEX-WRF runs over WA are available. The large spread in projected rainfall for different forcing data and/or regional models suggests to increase the ensemble size with further combinations of GCMs and RCMs for a better estimation of the projected uncertainty. With a higher resolution in time and space, our data can serve as input for further downscaling experiments at convection-resolving resolution (wascal2wps code available).